

# Chapter 1

## Mutant and Transgenic Zebrafish in Modeling Neurobehavioral Disorders

Carisa L. Bergner, Rupert J. Egan, Peter C. Hart, Jonathan M. Cachat, Peter R. Canavello, and Allan V. Kalueff

### Abstract

Zebrafish have traditionally been used as effective genetic and developmental models in biomedical research. Recently, the scope and utility of zebrafish in biomedical research has been further expanded with the implementation of new genetic techniques aimed at developing translational models of human pathogenesis. Additionally, screens measuring specific neurobehavioral and developmental phenotypes have proven to be very robust. This chapter further discusses the utility of zebrafish in biomedical research and highlights some of the genetic techniques used in the creation of transgenic and mutant strains. Behavioral phenotypes of genetically altered zebrafish are also discussed with respect to both their robust stress responses and similarity to human disorders. Specific emphasis is placed on human brain pathogenesis and neurodevelopmental abnormalities, especially as they relate to stress and anxiety spectrum disorders.

**Key words:** Zebrafish, Genetics, Translational models

---

### 1. The Utility of Zebrafish as a Model in Biomedical Research

Biomedical research relies on animal models to elucidate the cellular and molecular basis of human pathogenesis in order to generate efficacious treatments. While many mammalian species have been utilized due to their evolutionary proximity and homology to humans, zebrafish are quickly becoming a valuable model in many areas of research (1).

Zebrafish (*danio rerio*) have been traditionally used to explore the genetic and physiological mechanisms involved in the functioning of a variety of systems, including the cardiovascular (2), immune (3), and nervous systems (4). Additionally, the optical

transparency of the zebrafish has been widely exploited in research on the effects of genetic manipulation on a system's (e.g., nervous, circulatory, endocrine) development and function (5,6). Ease of genetic manipulation, high fecundity, and quick development have made the zebrafish model a useful tool for screening and investigating the genetic factors involved in the disposition to and progression of pathogenesis (1). The combination of transparent embryo development and the ease of embryological manipulations has fostered numerous advances in developmental biology, by allowing direct *in vivo* observation of cellular events (1). These techniques (cloning, mutagenesis, transgenesis, and mapping approaches) when applied to zebrafish allow researchers to use invertebrate-style forward genetics on a vertebrate organism (1). As a model of disease, zebrafish offer numerous advantages over mammalian models despite differences in anatomy. The ability to investigate the onset and progression of pathological processes in a suitable timeframe and *in vivo* is particularly notable. Contemporary research also demonstrates the robustness of behavioral phenotypes displayed by zebrafish in response to genetic manipulation and pharmacological treatment.

There are also drawbacks to the use of zebrafish in biomedical research. Zebrafish have developed a duplicate genome, although not all duplicated genes have been retained through time (7). Therefore, zebrafish possess two genes for many mouse orthologs, although the exact number of preserved duplicate genes remains unknown. It is frequently argued that further comprehension of zebrafish gene function will only uncover invalid redundant and species-specific information (7). However, duplicate genes can also provide significant advantages, particularly in relation to sub-functionalisation. In this technique, zebrafish co-orthologs represent selected expression patterns and developmental functions of mouse orthologs (7). In this way, restricted expression of zebrafish genes, in comparison to the corresponding mouse orthologs, may lead to an improved comprehension of developmental relations in cell lineage or tissue patterning in mice (7).

---

## 2. Zebrafish as Models of Brain Disorders

A zebrafish experimental model of stress and anxiety has recently been developed to investigate the physiological mechanisms and neural biomarkers associated with anxiety-like phenotypes. When the zebrafish is exposed to a novel environment, it dives to the bottom and gradually explores as it habituates to the test apparatus. Inhibited exploratory movement, reduced speed, and increased frequency of escape behaviors are associated with higher

levels of anxiety elicited by stressors (e.g., novel environment, predator exposure) (8–10). These behaviors are highly sensitive to pharmacological treatment, as exploratory behavior is markedly elevated in response to treatment with anxiolytic drug, including benzodiazepines, SSRIs, nicotine, and ethanol (9,11–13). Conversely, chronic stressful stimuli (e.g., predator exposure, alarm pheromone), anxiogenic drugs, and drug withdrawal have been shown to increase anxiety-like behavior in this paradigm, as zebrafish show latency to explore the upper layer of the novel environment, spend significantly less time exploring, exhibit more erratic movements, and freeze often (10,12,14). This simple yet high throughput test can be used as an approach in quickly and accurately identifying the biomarkers linked to a disorder and in screening the efficacy of different pharmacological treatments.

Genomic synteny and resulting cellular morphology and neurophysiology are largely conserved in zebrafish (15), supporting the translatability of data to advance a research on psychiatric disorders in humans. Transgenic zebrafish show much promise for developing high throughput animal models of complex brain disorders, such as Huntington's, Alzheimer's, and possibly schizophrenia (16–18). The common application of transgenic zebrafish models has been as a screening assay to identify the genetic and physiological mechanisms involved in brain disorders; however, development of behavioral models may provide further insight towards the diagnosis and pharmacological treatment of these disorders. Research using the zebrafish model of stress and anxiety must incorporate transgenic zebrafish models of brain disorders, to correctly assess the genetic factors that precipitate abnormal neurobiological, physiological, and behavioral phenotypes.

---

### 3. The Generation of a Transgenic or Mutant Zebrafish

The most frequently utilized methods of transgenic line formation include plasmid DNA or bacterial artificial chromosome (BAC) injections into the cytoplasm of one-cell stage embryos (19). However, these procedures yield germline transgenics with just 5–20% efficiency (19). Therefore, transposon-mediated systems become increasingly popular, due to their higher approximated efficiency of 50–80% (1). This method is rapid, effective, and produces lines that facilitate genetic screening. Other advantages of transposon-mediated transgenesis include vectors that are more simply engineered than in other methods, and the potential for the remobilization of existing insertions to generate new insertions (19).

Transposon-mediated transgenesis relies on transposon vectors, which use the *cis*-acting elements from naturally occurring transposons in juxtaposition with the gene generally encoded by the transposon in *trans* (19). It is this gene that encodes transposase, the catalyzing enzyme in both the excision of the vector from plasmid, and the assimilation into the host genome (19). To express a gene of interest or create a gene trap, researchers may engineer specific sequences into the vector rather than utilize the transposase gene (19). Plasmids containing the transposon vector and the mRNA, which encodes the transposase, are coinjected into embryos at the one-cell stage (19). The injected embryos will be mosaic for integration events, due to rapid divisions during the early stages of zebrafish development.

Although transgenic zebrafish may be created through the use of transposons, or the injection of plasmid DNA or pseudotyped retroviruses, retroviruses are the only efficient insertional elements in the generation of mutant zebrafish (19). This insertional mutagenesis allows for the mutated gene to be easily identified with the help of a molecular tag at the mutagenic lesion site (19). Sequencing of DNA flanking the insertion has been shown to be sufficient in the identification of the mutant gene in roughly 80% of mutants (19).

Retroviral-mediated insertional mutagenesis requires the injection of a virus into embryos between the 512 and 2,000-cell stage (19). These embryos will then grow into founder fish (F0), which have mosaic germ lines and upwards of 25–30 different insertions (Amsterdam). Quality control assays are typically performed on viral stocks and founder injections before breeding founders. F1 fish, however, are not mosaic, and transfer their insertions in a Mendelian manner (19). When F1 pools reach 6 weeks of age, tail DNA can be isolated from 30 fish in order to identify subjects with the most insertions. F1 zebrafish with the most insertions may then be crossed with other multiple insertion fish to produce an F2 pool. At minimum, 6 crosses are required to screen at the F2 level, resulting in almost every insertion being homozygosed in at least one of the crosses (19).

In addition to retroviral methods, chemical mutagens may also create random mutagenesis. Zebrafish mutants can be generated through the exposure to ethylnitrosourea (ENU), a popular mutagen that typically produces point mutations (1). Due to the relative resistance of zebrafish to ENU toxicity, higher levels of mutagenesis can be achieved and defined loci can be affected with greater accuracy than in the other vertebrates (1). Chemical mutagenesis generally supersedes retroviral mutagenesis in efficiency.

#### 4. Genetically Modified Zebrafish Express Marked Behavioral Phenotypes

Although zebrafish models of stress and anxiety have only recently been developed, research focusing on the effects of genetic manipulation on behavioral phenotypes is being conducted in conjunction with physiological analyses. Using forward and reverse genetic engineering, transgenic zebrafish strains have been established to model brain disorders and to screen the genetic factors that induce or intensify pathogenesis (19,20). Typically, zebrafish models are used to analyze the physiological mechanisms disrupted by genetic mutation that lead to neurobiological morphology relevant to psychiatric disorders. However, recent research has also assessed the behavioral phenotypes associated with these neurological abnormalities.

Several transgenic zebrafish exhibit robust aberrant behavioral phenotypes linked to the knockout of specific target genes. For example, *nevermind* (*nev*) gene mutant zebrafish display severe disruption of optic nerve innervation (21). While muscle morphology is normal, *nev* dorsal retinotectal axon projections terminate on both the dorsal and ventral side of the tectum, resulting in atypical swimming phenotypes (22). Specifically, these mutants display a corkscrew swimming pattern, in which they rotate around their long body axis (22). Similarly, sphingosylphosphorylcholine (SPC, a lipid mediator) KO zebrafish perform spontaneous erratic movements and escape behaviors (e.g., rapid turning) without provocation from stressful stimuli (22).

Many of the transgenic zebrafish models in current use focus on abnormal developmental patterns that prevent proper innervations between nuclei and in turn disrupt neurophysiology. One example of this is the mutation of the *Lhx2* homolog, *bel*, a transcription factor involved in retinotectal axonal growth. The optic chiasm does not form in these zebrafish mutants resulting in a robust phenotype that simulates a related human disorder, non-decussating retinal-fugal fiber syndrome (NDRFF). In zebrafish, achiasmatic-induced oculomotor deficits generate spontaneous eye oscillations that may model congenital nystagmus in humans, in addition to causing reversed perception of visual stimuli, misappropriated eye movements, and circling swimming behavior. Research utilizing this strain may help investigate developmental factors and possible treatments in patients suffering from oculomotor and visual impairment (23).

Vertebrate optokinetic nystagmus (OKN) is a compensatory oculomotor behavior that is evoked by movement of the visual environment. Its primary function is to stabilize visual images on the retina. It is possible to evoke this response from many vertebrate species, including zebrafish, in an experimental environment.

This enables for the screening of mutations affecting visual system development. Rick and colleagues (24) have shown that *bella-donna* (*bel*) mutant larvae exhibit abnormal functioning of the OKN with about 40% displaying reversal of the OKN upon experimentally induced visual stimulation. The behavioral deficit observed in OKN function is caused by defective, non functional optic chiasm formation in the achiasmatic mutants displaying a reversal of OKN (*bel* mutants). This study highlights how a forward genetic approach can be used in the zebrafish to uncover very specific behavioral deficits. Further, the analysis of behavioral mutations in combination with neuroanatomical analysis links behavioral abnormalities with their underlying structural deficits. This combination approach of analyzing the neural circuits by genetic means allows the researchers to establish a structure–function relationships of morphological defects and elucidate the resulting behavioral consequences (24).

Studies employing genetic classification in conjunction with behavioral analysis have been used to further examine the defects in the visual system of zebrafish. In particular, the problems of retinal degeneration and abnormal visual system development have been examined. Behavioral assays, along with morphological, electrophysiological, cellular, and molecular methods have proven to be powerful tools for examining this complex system (25). Advances have been made in regards to the isolation and classification of zebrafish with visual system deficits (26). Indeed, behavioral screens have been developed with the ability to isolate specific disturbances in the visual processing system of zebrafish (27). Advances in our understanding of disorders related to visual system development and visual system processing in zebrafish may lead to novel treatments for human visual system disorders, and highlight the translatability of zebrafish as genetic models of human pathogenesis.

In another transgenic model, *nacre* zebrafish that lack melanophore stripes were used in conjunction with wild-type zebrafish to examine the effects of altered morphology on social behavior and mating (28). Analyzing the distance between subjects within a shoal and the preference to swim near morphologically similar zebrafish showed no selectivity in regard to visual characteristics. As in other fish, morphological alterations may affect behavioral responses in zebrafish. For example, pigment patterns have been found to be an important factor in how zebrafish perceive shoalmates (29). Interestingly, there are distinct sexual differences with male zebrafish preferences depending on the visibility and regularity of stripe patterns, while the factors underlying females preferences are unclear.

In a similar experiment, mutant strains of zebrafish possessing either truncated (*short-fin*) or elongated (*long-fin*, Fig. 1.1) fin genes were generated and social/sexual interactions were observed. While there was no preference of the WT or *short-fin*



Fig. 1.1. Wild type (*top*) and long-fin mutant (*bottom*) zebrafish.

female zebrafish, *long-fin* female zebrafish showed an increased social interaction with *long-fin* over WT male zebrafish. Since female zebrafish are reported to select mates based on body size rather than fin size, and because the *short-fin* mutant showed no preference, the *long-fin* mutation is thought to be involved in social perception (30).

Additionally, a zebrafish model of Huntington's disease (HD) was generated through a reverse genetics technique, using a morpholino (antisense morpholino-modified oligonucleotides) to inhibit translation of the protein huntingtin (16,31). The huntingtin (Htt) *knockdown* zebrafish displayed abnormal physiological phenotypes associated with HD in humans, including reduced expression of brain-derived neurotrophic factor (BDNF), widespread neuronal apoptosis, and other developmental disruptions linked to HD. Unfortunately, no behavioral testing was performed in this experiment; however, a spatial alternation screen using reward stimuli as described by Williams and colleagues (32) could illuminate the learning and memory deficits associated with HD. In addition, assessing anhedonia, motor coordination, and anxiety in Htt *knockdown* zebrafish may help in fully defining a zebrafish transgenic model of HD.

Expression patterns in genes related to deafness begin to provide a basis for understanding the roles of various cell types within the cochlea and bring to light the causes of both syndromic and non-syndromic forms of deafness (33). Recently, a large scale mutagenesis screen in zebrafish identified 8 genes that are essential for proper balance and hearing (33). Zebrafish have proven to be effective models with respect to studying the molecular basis

of development and function of sensory hair cells (33). Importantly, anatomical studies have shown that the structure of the vestibular inner ear is highly conserved between teleost fish and higher animals. Additionally, fish sensory hair cells are morphologically and physiologically similar to hair cells in higher vertebrates (33). As such, zebrafish emerge as a novel translational model for examining the pathogenesis that occurs as a result of mutations in deafness genes (33).

Zebrafish are also proving useful as genetic models to examine the complex processes involved in the development of addiction, which is a principle health issue throughout the world. The use of forward genetic manipulation in conjunction with a selective behavioral screen for the classification of cocaine sensitivity in mutant zebrafish has helped to identify candidate genes responsible for sensitivity to addiction. In the highlighted experiment, a cocaine-induced conditioned place preference developed in a sensitivity-dependent manner. Sensitivity to cocaine, and the resulting behavioral phenotypes observed in the conditioned place preference paradigm, was shown to be dependent on the dominant mutations in single genes (33). Indeed, certain mutagenized zebrafish exhibited abnormally low responses to cocaine. Performance profiles of these zebrafish in additional behavioral screens measuring visual adaptation and learning suggested that their low response to cocaine was the result of mutations in genes that affect dopaminergic signaling in both the retina and the brain (34). This forward genetics approach in which the genome is mutagenized, resulting phenotypes characterized through behavioral screening, and underlying genes classified and cloned using a vertebrate species represents a valuable tool to advance the research. Zebrafish in particular are an idea choice for the forward genetic manipulation due to their ability to reproduce quickly, and their low maintenance cost (34). These benefits along with the fact that zebrafish lend themselves well to behavioral observation paradigms further expands their role as translational models in biomedical research.

---

## 5. Conclusion

Zebrafish represent effective models for translational research, spanning the developmental, genetic, and behavioral level. Their utility within the field of biomedical research is unquestionable. However, in order to fully realize their scope of impact researchers must combine approaches in order to create effective models that exhibit translatable validity towards the treatment of human pathogenesis. The development of new genetic techniques and their application towards the creation of mutant and transgenic

lines of zebrafish, merged with proper behavioral phenotype classification of genetically modified and mutant zebrafish, will allow the researchers to parallel human disorders with much greater accuracy. These developments will lead to a better understanding of the complex nature of neurobehavioral and neurodevelopmental disorders. Through the combined efforts of biomedical researchers in all areas of research the development of novel, effective treatments for neurobehavioral and neurodevelopmental disorders will be achieved.

---

## Acknowledgments

Supported by the NARSAD YI Award, Georgetown University's Stress Physiology and Research Center (SPaRC) and Tulane University Intramural Research Funds.

## References

1. Lieschke GJ, Currie PD (2007) Animal models of human disease: zebrafish swim into view. *Nat Rev Genet* 8:353–367
2. Stainier DY, (2001) Zebrafish genetics and vertebrate heart formation. *Nat Rev Genet*, 2(1):39–48
3. Sullivan C, Kim CH (2008) Zebrafish as a model for infectious disease and immune function. *Fish Shellfish Immunol* 25:341–350
4. Pogoda HM, Hammerschmidt M (2007) Molecular genetics of pituitary development in zebrafish. *Semin Cell Dev Biol* 18:543–558
5. Dale RM, Sisson BE, Topczewski J (2009) The emerging role of Wnt/PCP signaling in organ formation. *Zebrafish* 6(1):9–14
6. Segner H (2009) Zebrafish (*Danio rerio*) as a model organism for investigating endocrine disruption. *Comp Biochem Physiol C Toxicol Pharmacol* 149:187–195
7. Key B, Devine CA (2003) Zebrafish as an experimental model: strategies for developmental and molecular neurobiology studies. *Methods Cell Sci* 25:1–6
8. Bass SL, Gerlai R (2008) Zebrafish (*Danio rerio*) responds differentially to stimulus fish: the effects of sympatric and allopatric predators and harmless fish. *Behav Brain Res* 186:107–117
9. Levin ED, Bencan Z, Cerutti DT (2007) Anxiolytic effects of nicotine in zebrafish. *Physiol Behav* 90:54–58
10. Speedie N, Gerlai R (2008) Alarm substance induced behavioral responses in zebrafish (*Danio rerio*). *Behav Brain Res* 188:168–177
11. Airhart MJ, Lee DH, Wilson TD, Miller BE, Miller MN, Skalko RG (2007) Movement disorders and neurochemical changes in zebrafish larvae after bath exposure to fluoxetine (PROZAC). *Neurotoxicol Teratol* 29:652–664
12. Gerlai R, Ahmad F, Prajapati S (2008) Differences in acute alcohol-induced behavioral responses among zebrafish populations. *Alcohol Clin Exp Res* 32:1763–1773
13. Gerlai R, Lee V, Blaser R (2006) Effects of acute and chronic ethanol exposure on the behavior of adult zebrafish (*Danio rerio*). *Pharmacol Biochem Behav* 85:752–761
14. Baraban SC, Taylor MR, Castro PA, Baier H (2005) Pentylentetrazole induced changes in zebrafish behavior, neural activity and c-fos expression. *Neuroscience* 131:759–768
15. Detrich HW 3rd, Yergeau DA (2004) Comparative genomics in erythropoietic gene discovery: synergisms between the Antarctic icefishes and the zebrafish. *Methods Cell Biol* 77:475–503
16. Diekmann H, Anichtchik O, Fleming A, Futter M, Goldsmith P, Roach A et al (2009) Decreased BDNF levels are a major contributor to the embryonic phenotype of huntingtin knock-down zebrafish. *J Neurosci* 29:1343–1349
17. Lee JA, Cole GJ (2007) Generation of transgenic zebrafish expressing green fluorescent protein under control of zebrafish amyloid precursor protein gene regulatory elements. *Zebrafish* 4:277–286

18. Wood JD, Bonath F, Kumar S, Ross CA, Cunliffe VT (2009) Disrupted-in-schizophrenia 1 and neuregulin 1 are required for the specification of oligodendrocytes and neurones in the zebrafish brain. *Hum Mol Genet* 18:391–404
19. Amsterdam A, Becker TS (2005) Transgenes as screening tools to probe and manipulate the zebrafish genome. *Dev Dyn* 234:255–268
20. Zon LI, Peterson RT (2005) In vivo drug discovery in the zebrafish. *Nat Rev Drug Discov* 4:35–44
21. Solnica-Krezel L (1999) Pattern formation in zebrafish—fruitful liaisons between embryology and genetics. *Curr Top Dev Biol* 41:1–35
22. Patton EE, Zon LI (2001) The art and design of genetic screens: zebrafish. *Nat Rev Genet* 2:956–966
23. Huang YY, Rinner O, Hedinger P, Liu SC, Neuhaus SC (2006) Oculomotor instabilities in zebrafish mutant *belladonna*: a behavioral model for congenital nystagmus caused by axonal misrouting. *J Neurosci* 26:9873–9880
24. Rick JM, Horschke I, Neuhaus SC (2000) Optokinetic behavior is reversed in achiasmatic mutant zebrafish larvae. *Curr Biol* 10:595–598
25. Li L (2001) Zebrafish mutants: behavioral genetic studies of visual system defects. *Dev Dyn* 221:365–372
26. Brockerhoff SE, Hurley JB, Janssen-Bienhold U, Neuhaus SC, Driever W, Dowling JE (1995) A behavioral screen for isolating zebrafish mutants with visual system defects. *Proc Natl Acad Sci USA* 92:10545–10549
27. Brockerhoff SE, Dowling JE, Hurley JB (1998) Zebrafish retinal mutants. *Vision Res* 38:1335–1339
29. Engezer RE, Barbiano LA, Ryan MJ, Parichy DM (2007) Timing and plasticity of shoaling behaviour in the zebrafish. *Danio rerio. Anim Behav* 74: 1269–1275
29. Engeszer RE, et al., (2008) Sex-specific perceptual spaces for a vertebrate basal social aggregative behavior. *Proc Natl Acad Sci USA*, 105(3):929–933
30. Gumm JM, Snekser JL, Iovine MK (2009) Fin-mutant female zebrafish (*Danio rerio*) exhibit differences in association preferences for male fin length. *Behav Processes* 80:35–38
31. Nasevicius A, Ekker SC (2000) Effective targeted gene ‘knockdown’ in zebrafish. *Nat Genet* 26:216–220
32. Williams FE, White D, Messer WS (2002) A simple spatial alternation task for assessing memory function in zebrafish. *Behav Processes* 58:125–132
33. Ernest S, Rauch GJ, Haffter P, Geisler R, Petit C, Nicolson T (2000) *Mariner* is defective in myosin VIIA: a zebrafish model for human hereditary deafness. *Hum Mol Genet* 9:2189–2196
34. Darland T, Dowling JE (2001) Behavioral screening for cocaine sensitivity in mutagenized zebrafish. *Proc Natl Acad Sci USA* 98:11691–11696